

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/009218

International filing date: 18 March 2005 (18.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/575,133  
Filing date: 28 May 2004 (28.05.2004)

Date of receipt at the International Bureau: 25 April 2005 (25.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1308296

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*April 13, 2005*

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.**

**APPLICATION NUMBER: 60/575,133**

**FILING DATE: May 28, 2004**

**RELATED PCT APPLICATION NUMBER: PCT/US05/09218**



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

Attorney Docket No.: 10620.010-US

PATENT

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

Mail Stop Provisional Patent Application  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Sir:

This is a request for filing a provisional application under 37 CFR 1.53(c).

INVENTOR(S)		
First Name	Middle Name	Family Name
1.Swapnil 2.Henrik 3.Henrik 4.Anders 5.Malcolm		Bhargava Bisgard-Frantzen Frisner Vikso-Nielsen Johal
TITLE OF THE INVENTION		
Liquefaction Process		

The following application parts are enclosed:

☒ specification 23 pages ☒ Sequence Listing 3 pages

☒ Abstract 1 page ☒ Drawings 1 pages


An application data sheet is enclosed.

Direct all correspondence to Customer Number 25908.

Please charge the required fee, estimated to be \$160, to Novozymes North America, Inc., Deposit Account No. 50-1701. A duplicate of this sheet is enclosed.

Respectfully submitted,

Date: May 28, 2004

  
\_\_\_\_\_  
Jason I. Garbell, Reg. No. 44,116  
Novozymes North America, Inc.  
500 Fifth Avenue, Suite 1600  
New York, NY 10110  
(212) 840-0097

Attorney Docket No.: 10620.010-US

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

EXPRESS MAIL CERTIFICATE

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Re: U.S. Provisional Application for  
"Liquefaction Process"  
Applicants: Bhargava et al.

Sir:

Express Mail Label No. EV 403903045 US

Date of Deposit May 28, 2004

I hereby certify that the following attached paper(s) or fee

1. Filing Under 37 C.F.R. §1.53(c) (in duplicate)
2. Provisional Application
3. Application Data Sheet
4. Sequence Listing

are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 C.F.R. 1.10 on the date indicated above and is addressed to the address indicated above.

Julie Tabarovsky

(Name of person mailing paper(s) or fee)

Julie Tabarovsky

(Signature of person mailing paper(s) or fee)

Mailing Address:

Novozymes North America, Inc.  
500 Fifth Avenue, Suite 1600  
New York, NY 10110  
(212) 840-0097

## **LIQUEFACTION PROCES**

### **FIELD OF THE INVENTION**

The present invention relates to an improved method of liquefying starch-containing material suitable as a step in processes for producing syrups and fermentation products, such as especially ethanol. The invention also relates to processes for producing ethanol comprising liquefying starch-containing starting material in accordance with the liquefaction method of the invention.

### **BACKGROUND OF THE INVENTION**

Liquefaction is a well known process step in the art of producing syrups and fermentation products, such as ethanol, from starch-containing materials. During liquefaction starch is converted to shorter chains and less viscous dextrans. Generally liquefaction involves gelatinization of starch simultaneously with or followed by addition of alpha-amylase.

WO 02/38787 (Novozymes) disclose a method of producing ethanol by fermentation comprising carrying out secondary liquefaction in the presence of a thermostable acid alpha-amylase or a thermostable maltogenic acid alpha-amylase.

Even though liquefaction has already been improved significantly there is still a need for improving liquefaction suitable for in syrups and fermentation product producing processes.

### **SUMMARY OF THE INVENTION**

The object of the present invention is to provide an improved method of liquefying starch-containing material suitable as a step in processes for producing syrups and fermentation products, such as especially ethanol. The invention also provides an ethanol producing process including a liquefaction method of the invention.

The present inventors have found that when liquefaction is carried out on starch-containing material according to the present invention a number of advantages are obtained. For instance, the inventors have showed that a DE above 20 may be obtained without using more enzyme than corresponding prior art processes which reaches a DE of around 12. Further, reduced viscosity was observed. This eases handling of the liquefied material and reduces the cost of pumping the liquefied material to down stream process equipment such as a fermentor. Furthermore, the enzyme cost is also reduced. It was also found that the sugar profile of the liquefied mash had a decreased DP<sub>4+</sub> content and increased DP<sub>1-3</sub> content compared to corresponding prior art methods using higher amounts of enzyme. The higher DP<sub>1-3</sub> content makes the liquefied mash easier and potentially faster to ferment by the yeast during fermentation, e.g.,

an ethanol fermentation process. It could be attributed to the fact that small sugars are released pertaining to the (acid) alpha-amylase action. These small sugars, e.g., glucose, maltose, maltotriose can be directly metabolized by yeast and therefore making SSF more effective and faster.

Further, also the residual DP<sub>4+</sub> content were after fermentation found to be higher than in corresponding prior art processes. This indicates a better utilization of the starch-containing starting material.

The abbreviation "DE" stands for "Dextrose equivalents" and is a measure of reducing ends on C<sub>6</sub> carbohydrates. Pure dextrose (glucose) has DE of 100. Dextrose is a reducing sugar. Whenever an amylase hydrolyzes a glucose-glucose bond in starch, two new glucose end-groups are exposed. At least one of these can act as a reducing sugar. Therefore the degree of hydrolysis can be measured as an increase in reducing sugars. The value obtained is compared to a standard curve based on pure glucose - hence the term dextrose equivalent. In other words: DE (dextrose equivalent) is defined as the amount of reducing carbohydrate (measured as dextrose-equivalents) in a sample expressed as w/w% of the total amount of dissolved dry matter.

According to the first aspect the invention relates to a method of liquefying starch-containing material, wherein the method comprises the steps of:

(a) treating the starch-containing material with a bacterial alpha-amylase at a temperature around 70-90°C for 15-90 minutes,

(b) treating the material obtained in step (a) with an alpha-amylase at a temperature between 60-80°C for 30-90 minutes.

The term "mash" is used for liquefied starch-containing material, such as liquefied whole grain.

In one embodiment of the invention the starch-containing material is jet-cooking at 90-120°C, preferably around 105°C, for 1-15 minutes, preferably for 3-10 minute, especially around 5 minutes, before step (a).

The mash has after the secondary liquefaction in step (b) a DE value above 16, preferably above 18, especially above 20, such as a DE value in the range from 16 to 30, preferably in the range from 18 to 25.

In a second aspect the invention provides a process of producing ethanol from starch-containing material by fermentation, said process comprises the steps of:

(i) liquefying starch-containing material according to the liquefaction method of the invention;

- (ii) saccharifying the liquefied mash obtained;
- (iii) fermenting the material obtained in step (ii).

Optionally the ethanol is recovery after fermentation. In an embodiment the saccharification and fermentation is carried out as a simultaneous saccharification and fermentation process (SSF process).

## BRIEF DESCRIPTION OF THE INVENTION

**Fig. 1:** Ethanol yields for six liquefaction treatments for 0.3 AGU/g DS of Glucoamylase SF liquefaction treatments

## DESCRIPTION OF THE INVENTION

The present invention provides an improved liquefaction process suitable as a step in processes for producing fermentation products such as especially ethanol. The invention also relates to a process of producing ethanol comprising the liquefaction method of the invention. Where the end product is ethanol it may be used as, e.g., fuel ethanol; drinking ethanol, i.e., potable neutral spirits; or industrial ethanol.

### Liquefaction

According to the present invention "liquefaction" is a process step in which starch-containing material, preferably milled (whole) grain, is broken down (hydrolyzed) into maltodextrins (dextrins).

Initially an aqueous slurry containing preferably from 10-40 wt-%, especially 25-35 wt-% starch-containing material is prepared. The starch-containing material is preferably milled whole grain. Then the starch-containing material is incubated with a bacterial alpha-amylase, preferably one or more *Bacillus* alpha-amylase(s), and may in one embodiment be followed by a jet-cooking step carried out between 90-120°C, preferably around 105°C, for 1-15 minutes, preferably for 3-10 minutes, especially around 5 minutes, to complete gelatinization of the slurry. However, it is to be understood that the method of the invention may also be carried out without a jet-cooking step. After incubation with bacterial alpha-amylase, with or without jet-cooking, the temperature is adjusted to 60-80°C and the material is incubated for 30 to 90 minutes in the presence of an alpha-amylase, preferably an acid alpha-amylase, especially a fungal acid alpha-amylase, to finalize hydrolysis (secondary liquefaction).

Consequently, in the first aspect the invention provides a method for liquefying starch-containing material comprising the steps of:

(a) treating the starch-containing material with a bacterial alpha-amylase at a temperature around 70-90°C for 15-90 minutes,

(b) treating the material obtained in step (a) with an alpha-amylase at a temperature between 60-80°C for 30-90 minutes.

5           A liquefaction method of the invention is typically carried out at pH 4.5-6.5, in particular at a pH between 5 and 6.

          The alpha-amylase may be any alpha-amylase, preferred an alpha-amylase mentioned in the section "Alpha-amylases" below.

#### 10   Starch-containing material

          The starch-containing material used according to the present invention is selected from the group consisting of: tubers, roots and whole grain; and any combinations of the foregoing. In an embodiment, the starch-containing material is obtained from cereals. The starch-containing material may, e.g., be selected from the groups consisting of corns, cobs, wheat, barley, cas-  
15   sava, sorghum, rye, milo and potatoes; or any combination of the foregoing.

          If the liquefaction method of the invention is included in an ethanol process of the invention, the raw starch-containing material is preferably whole grain or at least mainly whole grain. A wide variety of starch-containing whole grain crops may be used as raw material including: corn (maize), milo, potato, cassava, sorghum, wheat, and barley. Thus, in one embodiment, the  
20   starch-containing material is whole grain selected from the group consisting of corn (maize), milo, potato, cassava, sorghum, wheat, and barley; or any combinations thereof. In a preferred embodiment, the starch-containing material is whole grain selected from the group consisting of corn, wheat and barley or any combinations thereof.

          The raw material may also consist of or comprise a side-stream from starch processing,  
25   e.g., C<sub>6</sub> carbohydrate containing process streams that are not suited for production of syrups.

#### Milling

          In a preferred embodiment of the invention the starch-containing material is milled before  
30   step (a), i.e., before the primary liquefaction. Thus, in a particular embodiment, the liquefaction method further comprises - prior to the primary liquefaction step (i.e., prior to step (a)), - the steps of:

- i.   milling of the starch-containing material, such as whole grain;
- ii.  forming a slurry comprising the milled starch-containing material and water.



The starch-containing material, such as whole grain, is milled in order to open up the structure and allowing for further processing. Two processes of milling are normally used in ethanol production processes: wet and dry milling. The term "dry milling" denotes milling of the whole grain. In dry milling the whole kernel is milled and used in the remaining part of the process. Wet milling gives a good separation of germ and meal (starch granules and protein) and is with a few exceptions applied at locations where there is a parallel production of syrups. Dry milling is preferred in processes aiming at producing ethanol.

The term "grinding" is also understood as milling. In a preferred embodiment of the invention dry milling is used

### **Ethanol process**

An ethanol production process of the invention generally involves the steps of liquefaction, saccharification, fermentation and optionally recovering the ethanol product, preferably by distillation.

According to this aspect, the invention relates to a process of producing ethanol from starch-containing material by fermentation, said method comprises the steps of: (i) liquefying said starch-containing material according to the liquefaction method of the invention;

(ii) saccharifying the liquefied mash obtained in step (i)

(iii) fermenting the material obtained in step (ii).

In an embodiment the saccharification and fermentation steps are carried out as a simultaneous saccharification and fermentation process (SSF process). In a preferred embodiment of the invention starch-containing raw material, such as whole grain, preferably corn, is dry milled in order to open up the structure and allow for further processing. The mash has before step (ii), i.e., after step (i), with or without jet-cooking, a DE value of above 16, preferably above 18, especially above 20, such as a DE value in the range from 16 to 30, preferably in the range from 18 to 25.

A specific embodiment of the ethanol process of the invention comprises the steps of;

1) liquefying starch-containing material in accordance with the liquefaction method of the invention;

2) liquefying the material obtained in step 1) in the presence of an alpha-amylase having an amino acid sequence which has at least 70% identity to SEQ ID NO:1; and

3) saccharifying the material obtained; and

4) fermenting to produce ethanol;

wherein the steps 1), 2), 3) and 4) is performed in the order 1), 2), 3), 4) or wherein 4) is performed simultaneously to or following 3).

In a preferred embodiment a jet-cooking step, as defined above, is included after step i).

In a preferred embodiment the alpha-amylase used in step ii) is at least 75%, 80%, 85% or at least 90%, e.g., at least 95%, 97%, 98%, or at least 99% identity to SEQ ID NO:1.

The mash has after step 2), with or without jet-cooking, a DE value of above 16, preferably above 18, especially above 20, such as a DE value in the range from 16 to 30, preferably in the range from 18 to 25.

## 10 Saccharification

"Saccharification" is a step in which the maltodextrin (such as, product from the liquefaction) is converted to low molecular sugars  $DP_{1-3}$  (i.e., carbohydrate source) that can be metabolized by a fermenting organism, such as, yeast. Saccharification steps are well known in the art and are typically performed enzymatically using at least a glucoamylase or one or more carbohydrate-source generating enzymes as will be defined below. The saccharification step comprised in the process for producing ethanol of the invention may be a well known saccharification step in the art. In one embodiment glucoamylase, alpha-glucosidase and/or acid alpha-amylase is used for treating the liquefied starch-containing material. A full saccharification step may last up to from 20 to 100 hours, preferably about 24 to about 72 hours, and is often carried out at temperatures from about 30 to 65°C, and at a pH between 4 and 6, normally around pH 4.5-5.0. However, it is often more preferred to do a pre-saccharification step, lasting for about 40 to 90 minutes, at temperature of between 30-65°C, typically about 60°C, followed by complete saccharification during fermentation in a simultaneous saccharification and fermentation process (SSF). The most widely used process in ethanol production is the simultaneous saccharification and fermentation (SSF) process, in which there is no holding stage for the saccharification, meaning that fermenting organism, such as yeast, and enzyme(s) is(are) added together. In SSF processes, it is common to introduce a pre-saccharification step at a temperature between 40 and 60°C, preferably around 50°C, just prior to the fermentation.

## 30 Fermentation

In an ethanol production process of the invention the fermenting organism is preferably yeast, which is applied to the saccharified material.

The term "fermenting organism" refers to any organism suitable for use in a desired fermentation process. Suitable fermenting organisms are according to the invention capable to ferment, i.e., convert sugars, such as glucose or maltose, directly or indirectly into ethanol. Ex-

amples of fermenting organisms include fungal organisms, such as yeast. Preferred yeast includes strains of the *Sacchromyces* spp., and in particular *Sacchromyces cerevisiae*. Commercially available yeast includes, e.g., RED STAR®/Lesaffre Ethanol Red (available from Red Star/Lesaffre, USA) FALI (available from Fleischmann's Yeast, a division of Burns Philp Food Inc., USA), SUPERSTART (available from Alltech), GERT STRAND (available from Gert Strand AB, Sweden) and FERMIOL (available from DSM Specialties). In preferred embodiments, yeast is applied to the saccharified mash. Fermentation is ongoing for 24-96 hours, such as typically 35-65 hours. In preferred embodiments, the temperature is generally between 26-34°C, in particular about 32°C, and the pH is generally from pH 3-6, preferably around pH 4-5. Yeast cells are preferably applied in amounts of  $10^5$  to  $10^{12}$ , preferably from  $10^7$  to  $10^{10}$ , especially  $5 \times 10^7$  viable yeast count per ml of fermentation broth. During the ethanol producing phase the yeast cell count should preferably be in the range from  $10^7$  to  $10^{10}$ , especially around  $2 \times 10^8$ . Further guidance in respect of using yeast for fermentation can be found in, e.g., "The alcohol Textbook" (Editors K. Jacques, T.P. Lyons and D.R.Kelsall, Nottingham University Press, United Kingdom 1999), which is hereby incorporated by reference.

#### Recovery of ethanol

Optionally the ethanol is recovered after fermentation, preferably by including the step of;

(iv) distillation to obtain the ethanol;

wherein the fermentation in step (iii) and the distillation in step (iv) is carried out simultaneously or separately/sequential; optionally followed by one or more process steps for further refinement of the ethanol.

#### Starch Conversion

The liquefaction method of the invention may also be included in a traditional starch conversion process for producing syrups such as glucose, maltose, malto-oligosaccharides and isomalto-oligosaccharides.

#### **Alpha-amylases**

According to the invention preferred alpha-amylases are of fungal or bacterial origin.

In an embodiment the bacterial alpha-amylase is a *Bacillus* alpha-amylase, such as, derived from a strain of *B. licheniformis*, *B. amyloliquefaciens*, and *B. stearothermophilus*. Other alpha-amylases include alpha-amylase derived from a strain of the *Bacillus* sp. NCIB 12289, NCIB 12512, NCIB 12513 or DSM 9375, all of which are described in detail in WO 95/26397,

and the alpha-amylase described by Tsukamoto et al., Biochemical and Biophysical Research Communications, 151 (1988), pp. 25-31.

Bacterial alpha-amylase variants and hybrids are described in WO 96/23873, WO 96/23874, WO 97/41213, and WO 99/19467.

5 Fungal alpha-amylases include alpha-amylases derived from a strain of *Aspergillus*, such as, *Aspergillus oryzae*, *Aspergillus niger*, or *A. kawashii* alpha-amylases. In a preferred embodiment, the alpha-amylase is an acid alpha-amylase. In a more preferred embodiment the acid alpha-amylase is an acid fungal alpha-amylase or an acid bacterial alpha-amylase. More preferably, the acid alpha-amylase is an acid fungal alpha-amylase derived from the genus *Aspergillus*. A commercially available acid fungal amylase is SP288 (available from Novozymes A/S, Denmark).

In an embodiment the alpha-amylase is an acid alpha-amylase. The term "acid alpha-amylase" means an alpha-amylase (E.C. 3.2.1.1) which added in an effective amount has activity at a pH in the range of 3.0 to 7.0, preferably from 3.5 to 6.0, or more preferably from 4.0-5.0.

15 A preferred acid fungal alpha-amylase is a Fungamyl-like alpha-amylase. In the present disclosure, the term "Fungamyl-like alpha-amylase" indicates an alpha-amylase which exhibits a high identity, i.e., more than 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85% 90%, 95 or even 99% identical to the amino acid sequence shown in SEQ ID NO: 10 in WO 96/23874. When used as a maltose generating enzyme fungal alpha-amylases may be added in an amount of 0.001-1.0 AFAU/g DS, preferably from 0.002-0.5 AFAU/g DS, preferably 0.02-0.1 AFAU/g DS.

20 Preferably the alpha-amylase is an acid alpha-amylase, preferably from the genus *Aspergillus*, preferably of the species *Aspergillus niger*. In a preferred embodiment the acid fungal alpha-amylase is the one from *A. niger* disclosed as "AMYA\_ASPNG" in the Swiss-prot/TrEMBL database under the primary accession no. P56271. Also variants of set acid fungal amylase having at least 70% identity, such as at least 80% or even at least 90%, 95%, 96%, 97%, 98% or 99% identity thereto is contemplated. In an embodiment the acid fungal alpha-amylase is the one disclosed in SEQ ID NO: 1, or a sequence being at least 70% identical, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99% identity to SEQ ID NO:1.

30 For purposes of the present invention, the degree of identity between two amino acid sequences is determined by the Clustal method (Higgins, 1989, CABIOS 5: 151-153) using the LASERGENE™ MEGALIGN™ software (DNASTAR, Inc., Madison, WI) with an identity table and the following multiple alignment parameters: Gap penalty of 10, and gap length penalty of 10. Pairwise alignment parameters were Ktuple=1, gap penalty=3, windows=5, and diagonals=5].

A preferred acid alpha-amylase for use in the present invention may be derived from a strain of *B. licheniformis*, *B. amyloliquefaciens*, and *B. stearothermophilus*. The alpha-amylase may also be an alpha-amylase variant. Specifically contemplated alpha-amylase variants are disclosed in US patent nos. 6,093,562, 6,297,038 or US patent no. 6,187,576 (hereby incorporated by reference).

Preferred commercial compositions comprising an alpha-amylase include MYCOLASE™ from DSM; BAN™, TERMAMYL™ SC, FUNGAMYL™, LIQUOZYME™ X and SAN™ SUPER, SAN™ EXTRA L from Novozymes A/S, Denmark) and CLARASE™ L-40,000, DEX-LO™, SPEYME FRED, SPEZYME™ AA, and SPEZYME™ DELTA AA (Genencor Int., USA), and the acid fungal alpha-amylase sold under the trade name SP 288 (available from Novozymes A/S, Denmark).

The bacterial alpha-amylase may be added in amounts as are well-known in the art. When measured in KNU units the alpha-amylase activity is preferably present in an amount of 0.5-5,000 NU/g of DS, in an amount of 1-500 AAU/kg of DS, or more preferably in an amount of 5-1,000 KNU/kg of DS, such as 10-100 KNU/kg DS.

Fungal acid alpha-amylase are preferably added in an amount of 0.001-10 AFAU/g of DS, in an amount of 0.01-0.25 AFAU/g of DS, or more preferably in an amount of 0.05-0.20 AFAU/kg of DS, such as around 0.1 AFAU/kg DS.

## **Carbohydrate-Source Generating Enzyme**

The term "carbohydrate-source generating enzyme" includes glucoamylase (being glucose generators), beta-amylase and maltogenic amylases (being maltose generators). A carbohydrate-source generating enzyme is capable of providing energy to the fermenting organism(s) used in a process of the invention for producing ethanol and/or may be converting directly or indirectly to a desired fermentation product, preferably ethanol. The carbohydrate-source generating enzyme may be mixtures of enzymes falling within the definition. Especially contemplated mixtures are mixtures of at least a glucoamylase and an alpha-amylase, especially an acid amylase, even more preferred an acid fungal alpha-amylase. The ratio between acidic fungal alpha-amylase activity (AFAU) per glucoamylase activity (AGU) (AFAU per AGU) may in an embodiment of the invention be at least 0.1, in particular at least 0.16, such as in the range from 0.12 to 0.50.

Examples of contemplated glucoamylases, maltogenic amylases, and beta-amylases are set forth in the sections above and below.

## Glucoamylase

A glucoamylase used according to the invention may be derived from any suitable source, e.g., derived from a microorganism or a plant. Preferred glucoamylases are of fungal or bacterial origin, selected from the group consisting of *Aspergillus* glucoamylases, in particular *A. niger* G1 or G2 glucoamylase (Boel et al. (1984), EMBO J. 3 (5), p. 1097-1102), or variants thereof, such as disclosed in WO 92/00381, WO 00/04136 add WO 01/04273 (from Novozymes, Denmark); the *A. awamori* glucoamylase (WO 84/02921), *A. oryzae* (Agric. Biol. Chem. (1991), 55 (4), p. 941-949), or variants or fragments thereof.

Other *Aspergillus* glucoamylase variants include variants to enhance the thermal stability: G137A and G139A (Chen et al. (1996), Prot. Eng. 9, 499-505); D257E and D293E/Q (Chen et al. (1995), Prot. Engng. 8, 575-582); N182 (Chen et al. (1994), Biochem. J. 301, 275-281); disulphide bonds, A246C (Fierobe et al. (1996), Biochemistry, 35, 8698-8704; and introduction of Pro residues in position A435 and S436 (Li et al. (1997), Protein Engng. 10, 1199-1204. Other glucoamylases include *Athelia rolfsii* (previously denoted *Corticium rolfsii*) glucoamylase (see US patent no. 4,727,026 and (Nagasaka,Y. et al. (1998) Purification and properties of the raw-starch-degrading glucoamylases from *Corticium rolfsii*, Appl Microbiol Biotechnol 50:323-330), *Talaromyces* glucoamylases, in particular, derived from *Talaromyces emersonii* (WO 99/28448), *Talaromyces leycettanus* (US patent no. Re. 32,153), *Talaromyces duponti*, *Talaromyces thermophilus* (US patent no. 4,587,215). Bacterial glucoamylases contemplated include glucoamylases from the genus *Clostridium*, in particular *C. thermoamylolyticum* (EP 135,138), and *C. thermohydrosulfuricum* (WO 86/01831).

Commercially available compositions comprising glucoamylase include AMG 200L; AMG 300 L; SAN™ SUPER, SAN™ EXTRA L, SPIRIZYME™ PLUS, SPIRIZYME™ FUEL, SPIRIZYME™ B4U and AMG™ E (from Novozymes A/S); OPTIDEX™ 300 (from Genencor Int.); AMIGASE™ and AMIGASE™ PLUS (from DSM); G-ZYME™ G900, G-ZYME™ and G990 ZR (from Genencor Int.).

Glucoamylases may in an embodiment be added in an amount of 0.02-20 AGU/g DS, preferably 0.1-10 AGU/g DS, such as 2 AGU/g DS.

## Beta-amylase

At least according to the invention the a beta-amylase (E.C 3.2.1.2) is the name traditionally given to exo-acting maltogenic amylases, which catalyze the hydrolysis of 1,4-alpha-glucosidic linkages in amylose, amylopectin and related glucose polymers. Maltose units are successively removed from the non-reducing chain ends in a step-wise manner until the mole-

cule is degraded or, in the case of amylopectin, until a branch point is reached. The maltose released has the beta anomeric configuration, hence the name beta-amylase.

Beta-amylases have been isolated from various plants and microorganisms (W.M. Fogarty and C.T. Kelly, Progress in Industrial Microbiology, vol. 15, pp. 112-115, 1979). These beta-amylases are characterized by having optimum temperatures in the range from 40°C to 65°C and optimum pH in the range from 4.5 to 7. A commercially available beta-amylase from barley is NOVOZYM™ WBA from Novozymes A/S, Denmark and SPEZYME™ BBA 1500 from Genencor Int., USA.

#### Maltogenic amylase

The amylase may also be a maltogenic alpha-amylase. A "maltogenic alpha-amylase" (glucan 1,4-alpha-maltohydrolase, E.C. 3.2.1.133) is able to hydrolyze amylose and amylopectin to maltose in the alpha-configuration. A maltogenic alpha-amylase from *Bacillus stearothermophilus* strain NCIB 11837 is commercially available from Novozymes A/S under the tradename MALTOGENASE™. Maltogenic alpha-amylases are described in US Patent nos. 4,598,048, 4,604,355 and 6,162,628, which are hereby incorporated by reference.

The maltogenic amylase may in a preferred embodiment be added in an amount of 0.05-5 mg total protein/gram DS or 0.05- 5 MANU/g DS.

#### **Production of Enzymes**

The enzymes referenced herein may be derived or obtained from any suitable origin, including, bacterial, fungal, yeast or mammalian origin. The term "derived" or means in this context that the enzyme may have been isolated from an organism where it is present natively, i.e., the identity of the amino acid sequence of the enzyme are identical to a native enzyme. The term "derived" also means that the enzymes may have been produced recombinantly in a host organism, the recombinant produced enzyme having either an identity identical to a native enzyme or having a modified amino acid sequence, e.g., having one or more amino acids which are deleted, inserted and/or substituted, i.e., a recombinantly produced enzyme which is a mutant and/or a fragment of a native amino acid sequence or an enzyme produced by nucleic acid shuffling processes known in the art. Within the meaning of a native enzyme are included natural variants. Furthermore, the term "derived" includes enzymes produced synthetically by, e.g., peptide synthesis. The term "derived" also encompasses enzymes which have been modified e.g., by glycosylation, phosphorylation, or by other chemical modification, whether *in vivo* or *in vitro*. The term "obtained" in this context means that the enzyme has an amino acid sequence identical to a native enzyme. The term encompasses an enzyme that has been isolated from an

organism where it is present natively, or one in which it has been expressed recombinantly in the same type of organism or another, or enzymes produced synthetically by, e.g., peptide synthesis. With respect to recombinantly produced enzymes the terms "obtained" and "derived" refers to the identity of the enzyme and not the identity of the host organism in which it is produced recombinantly.

The enzymes may also be purified. The term "purified" as used herein covers enzymes free from other components from the organism from which it is derived. The term "purified" also covers enzymes free from components from the native organism from which it is obtained. The enzymes may be purified, with only minor amounts of other proteins being present. The expression "other proteins" relate in particular to other enzymes. The term "purified" as used herein also refers to removal of other components, particularly other proteins and most particularly other enzymes present in the cell of origin of the enzyme of the invention. The enzyme may be "substantially pure," that is, free from other components from the organism in which it is produced, that is, for example, a host organism for recombinantly produced enzymes. In preferred embodiment, the enzymes are at least 75% (w/w) pure, more preferably at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% pure. In another preferred embodiment, the enzyme is 100% pure.

The enzymes used according to the present invention may be in any form suitable for use in the processes described herein, such as, e.g., in the form of a dry powder or granulate, a non-dusting granulate, a liquid, a stabilized liquid, or a protected enzyme. Granulates may be produced, e.g., as disclosed in US Patent Nos. 4,106,991 and US 4,661,452, and may optionally be coated by process known in the art. Liquid enzyme preparations may, for instance, be stabilized by adding stabilizers such as a sugar, a sugar alcohol or another polyol, lactic acid or another organic acid according to established process. Protected enzymes may be prepared according to the process disclosed in EP 238,216.

Even if not specifically mentioned in context of a method or process of the invention, it is to be understood that the enzyme(s) or agent(s) is(are) used in an "effective amount".

## **MATERIALS AND METHODS**

### Enzymes:

Bacterial Alpha-amylase A: *Bacillus stearothermophilus* alpha-amylase variant with the mutations: I181\*+G182\*+N193F disclosed in US patent no. 6,187,576 and available on request from Novozymes A/S, Denmark.



Fungal acid alpha-amylase B: *Aspergillus niger* alpha-amylase disclosed in SEQ ID NO: 1 and available from Novozymes A/S.

Glucoamylase T: Glucoamylase derived from *Talaromyces emersonii* and disclosed as SEQ ID NO: 7 in WO 99/28448.

Glucoamylase SF: Balanced blend of *Aspergillus niger* glucoamylase and *A. niger acid alpha-amylase* having a ratio between AGU and AFAU of approx. 9:1.

Stock solution for iodine method:

0.1N I<sub>2</sub>

· dissolve 1.3 g I<sub>2</sub> and 2.0 g KI into 100 mL DI water

**Methods:**

Alpha-amylase activity (KNU)

The amylolytic activity may be determined using potato starch as substrate. This method is based on the break-down of modified potato starch by the enzyme, and the reaction is followed by mixing samples of the starch/enzyme solution with an iodine solution. Initially, a blackish-blue color is formed, but during the break-down of the starch the blue color gets weaker and gradually turns into a reddish-brown, which is compared to a colored glass standard.

One Kilo Novo alpha amylase Unit (KNU) is defined as the amount of enzyme which, under standard conditions (i.e., at 37°C +/- 0.05; 0.0003 M Ca<sup>2+</sup>; and pH 5.6) dextrinizes 5260 mg starch dry substance Merck Amylum soluble.

A folder EB-SM-0009.02/01 describing this analytical method in more detail is available upon request to Novozymes A/S, Denmark, which folder is hereby included by reference.

Determination of FAU activity

One Fungal Alpha-Amylase Unit (FAU) is defined as the amount of enzyme, which breaks down 5.26 g starch (Merck Amylum soluble Erg. B.6, Batch 9947275) per hour based upon the following standard conditions:

Substrate . . . . . Soluble starch

Temperature . . . . . 37°C

pH. . . . . 4.7

Reaction time . . . . 7-20 minutes

Determination of acid alpha-amylase activity (AFAU)

5        Acid alpha-amylase activity is measured in AFAU (Acid Fungal Alpha-amylase Units), which are determined relative to an enzyme standard.

          The standard used is AMG 300 L (from Novozymes A/S, Denmark, glucoamylase wild-type *Aspergillus niger* G1, also disclosed in Boel et al. (1984), EMBO J. 3 (5), p. 1097-1102) and WO 92/00381). The neutral alpha-amylase in this AMG falls after storage at room  
10    temperature for 3 weeks from approx. 1 FAU/mL to below 0.05 FAU/mL.

          The acid alpha-amylase activity in this AMG standard is determined in accordance with the following description. In this method, 1 AFAU is defined as the amount of enzyme, which degrades 5.260 mg starch dry matter per hour under standard conditions.

          Iodine forms a blue complex with starch but not with its degradation products. The in-  
15    tensity of color is therefore directly proportional to the concentration of starch. Amylase activity is determined using reverse colorimetry as a reduction in the concentration of starch under specified analytic conditions.

	Alpha-amylase	
Starch + Iodine	→	Dextrins + Oligosaccharides
	40°C, pH 2.5	
Blue/violet	t=23 sec.	Decoloration

Standard conditions/reaction conditions: (per minute)

Substrate:	Starch, approx. 0.17 g/L
Buffer:	Citrate, approx. 0.03 M
Iodine (I <sub>2</sub> ):	0.03 g/L
CaCl <sub>2</sub> :	1.85 mM
pH:	2.50 ± 0.05
Incubation temperature:	40°C
Reaction time:	23 seconds
Wavelength:	lambda=590nm
Enzyme concentration:	0.025 AFAU/mL
Enzyme working range:	0.01-0.04 AFAU/mL

          If further details are preferred these can be found in EB-SM-0259.02/01 avail-  
20    able on request from Novozymes A/S, Denmark, and incorporated by reference.

### Acid Alpha-amylase Units (AAU)

The acid alpha-amylase activity can be measured in AAU (Acid Alpha-amylase Units), which is an absolute method. One Acid Amylase Unit (AAU) is the quantity of enzyme converting 1 g of starch (100% of dry matter) per hour under standardized conditions into a product having a transmission at 620 nm after reaction with an iodine solution of known strength equal to the one of a color reference.

Standard conditions/reaction conditions:

Substrate: Soluble starch. Concentration approx. 20 g DS/L.

Buffer: Citrate, approx. 0.13 M, pH=4.2

Iodine solution: 40.176 g potassium iodide + 0.088 g iodine/L

City water 15°-20°dH (German degree hardness)

pH: 4.2

Incubation temperature: 30°C

Reaction time: 11 minutes

Wavelength: 620nm

Enzyme concentration: 0.13-0.19 AAU/mL

Enzyme working range: 0.13-0.19 AAU/mL

The starch should be Lintner starch, which is a thin-boiling starch used in the laboratory as colorimetric indicator. Lintner starch is obtained by dilute hydrochloric acid treatment of native starch so that it retains the ability to color blue with iodine. Further details can be found in EP0140410B2, which disclosure is hereby included by reference.

### Glucoamylase activity (AGI)

Glucoamylase (equivalent to amyloglucosidase) converts starch into glucose. The amount of glucose is determined here by the glucose oxidase method for the activity determination. The method described in the section 76-11 Starch—Glucoamylase Method with Subsequent Measurement of Glucose with Glucose Oxidase in "Approved methods of the American Association of Cereal Chemists". Vol.1-2 AACC, from American Association of Cereal Chemists, (2000); ISBN: 1-891127-12-8.

One glucoamylase unit (AGI) is the quantity of enzyme which will form 1 micromol of glucose per minute under the standard conditions of the method.

Standard conditions/reaction conditions:

Substrate: Soluble starch.

Concentration approx. 16 g dry matter/L.

Buffer: Acetate, approx. 0.04 M, pH=4.3

pH: 4.3

Incubation temperature: 60°C

Reaction time: 15 minutes

Termination of the reaction: NaOH to a concentration of approximately 0.2 g/L (pH~9)

Enzyme concentration: 0.15-0.55 AAU/mL

The starch should be Lintner starch, which is a thin-boiling starch used in the laboratory as colorimetric indicator. Lintner starch is obtained by dilute hydrochloric acid treatment of native starch so that it retains the ability to color blue with iodine.

## 5 Glucoamylase activity (AGU)

The Novo Glucoamylase Unit (AGU) is defined as the amount of enzyme, which hydrolyzes 1 micromole maltose per minute under the standard conditions 37°C, pH 4.3, substrate: maltose 23.2 mM, buffer: acetate 0.1 M, reaction time 5 minutes.

10 An autoanalyzer system may be used. Mutarotase is added to the glucose dehydrogenase reagent so that any alpha-D-glucose present is turned into beta-D-glucose. Glucose dehydrogenase reacts specifically with beta-D-glucose in the reaction mentioned above, forming NADH which is determined using a photometer at 340 nm as a measure of the original glucose concentration.

AMG incubation:	
Substrate:	maltose 23.2 mM
Buffer:	acetate 0.1 M
pH:	4.30 ± 0.05
Incubation temperature:	37°C ± 1
Reaction time:	5 minutes
Enzyme working range:	0.5-4.0 AGU/mL

15

Color reaction:	
GlucDH:	430 U/L
Mutarotase:	9 U/L
NAD:	0.21 mM
Buffer:	phosphate 0.12 M; 0.15 M NaCl

pH:	7.60 ± 0.05
Incubation temperature:	37°C ± 1
Reaction time:	5 minutes
Wavelength:	340 nm

A folder (EB-SM-0131.02/01) describing this analytical method in more detail is available on request from Novozymes A/S, Denmark, which folder is hereby included by reference.

#### Determination of Maltogenic Amylase activity (MANU)

- 5 One MANU (Maltogenic Amylase Novo Unit) may be defined as the amount of enzyme required to release one micro mole of maltose per minute at a concentration of 10 mg of maltotriose (Sigma M 8378) substrate per ml of 0.1 M citrate buffer, pH 5.0 at 37°C for 30 minutes.

#### Standard iodine method

- 10 · boil small aliquot (10-20 mLs) of liquefied material in a test tube for several minutes  
 · cool in ice bath  
 · add 10-12 drops of the iodine solution  
 · mix and let sample sit in ice water for about 10 minutes

#### 15 Determination of Viscosity

The mash is heated to a temperature of 50-70°C, depending on the treatment. Following treatment viscosity is measured using a Haake VT02 rotation based viscosimeter. The unit of viscosity is centipois (cps), which is proportionally related to the viscosity level.

#### 20 Determination of DE (Dextrose Equivalent)

The DE value is measured using Fehlings liquid by forming a copper complex with the starch using pure glucose as a reference, which subsequently is quantified through iodometric titration. DE (dextrose equivalent) is defined as the amount of reducing carbohydrate (measured as dextrose-equivalents) in a sample expressed as w/w% of the total amount of dissolved dry matter. It may also be measured by the neocuproine assay (Dygert, Li Floridana(1965) Anal. Biochem. No 368). The principle of the neocuproine assay is that CuSO<sub>4</sub> is added to the sample, Cu<sup>2+</sup> is reduced by the reducing sugar and the formed neocuproine complex is measured at 450 nm.

#### **EXAMPLES**

30

## Example 1

### Liquefaction with bacterial and acid alpha amylase

400 mL of ground corn slurry is liquefied with 50 NU/g dry solids (DS) Bacterial Alpha-Amylase A. The corn mash has about 30% dry substance (pH 5.4). The mash is heated to 85°C and the viscosity and DE values are measured.

The mash is then treated with acid alpha-amylase B from *Aspergillus niger* having the amino acid sequence disclosed in SEQ ID NO:1. The enzyme loading is 0.10 AFAU/g dry solids. After 1.5 hours the viscosity and DE value are measured.

## Example 2

### The Effect of Alpha-Amylase Addition During Liquefaction on SSF Performance:

To investigate the effect of alpha-amylase (in form of Acid Alpha-Amylase B) addition in liquefaction, six different conditions for liquefaction were tested. To begin with ground corn was used to make 30% slurry with tap water. The pH in all the liquefactions were adjusted to 5.4 using diluted H<sub>2</sub>SO<sub>4</sub>. In the first two liquefactions (controls), Bacterial Alpha-Amylase A (50 NU/ g DS) was added and kept at 85°C for 1.5 and 4.5 hours, respectively. In the second set of runs (Acid Alpha-Amylase B test), the same process was followed, but incubation time with Bacterial Alpha-Amylase A at 85 °C was reduced to 0.5 hours, the temperature was then lowered and Acid Alpha-Amylase B was added (0.050 and 0.10 AFAU/g DS). The mixture was then kept at 70°C for 1 and 4 hours, respectively. Once the liquefaction was over, the reactions were stopped by adding 2 drops of HCl (4 N). Samples were withdrawn to analyze sugar profiles (using HPLC) and DE values. The liquefied samples were frozen and later subjected to SSF.

The effect of liquefaction treatment on SSF was evaluated via mini-scale fermentations. Samples after liquefaction were thawed and the pH was adjusted to 5.0 with diluted H<sub>2</sub>SO<sub>4</sub>. Two different dosages of Glucoamylase T derived from *Talaromyces emersonii* and Glucoamylase SF, respectively, were used for each treatment. Dosages used were 0.1 and 0.3 AGU/g DS. Approximately 4 grams of mash was added to 16 ml polystyrene tubes (Falcon 352025). Tubes were then dosed with the appropriate amount of Glucoamylase SF (0.3 AGU/g DS). Six replicates of each treatment were run. After dosing the tubes with enzyme, they were inoculated with 0.04 ml/g mash of yeast propagate that had been grown for 21 hours on corn mash. Vials were capped with a screw on lid which had been punctured with a very small needle to allow gas release and vortexed briefly before weighing and incubation at 32°C. Fermentation progress was followed by weighing the tubes over time. Tubes were vortexed briefly before each weighing. Weight loss values were converted to ethanol yield (g ethanol/g DS) by the following formula:

$$g_{ethanol} g DS = \frac{g_{CO_2} weight loss \times \frac{1 mol CO_2}{44.009 g CO_2} \times \frac{1 mole ethanol}{1 mol CO_2} \times \frac{46.094 g ethanol}{1 mole ethanol}}{g_{corn} in tube \times \% DS of corn}$$

- 5 Data from the liquefaction process shows that adding Acid Alpha-Amylase B in addition to Bacterial alpha-amylase A resulted in a significant increase in DE values.

Treatment	enzyme1	enzyme 2	DE
1	BAA(50)-85°C-1.5hr		7.27
2	BAA(50)-85°C-4.5hr		7.79
3	BAA(50)-85°C-0.5hr	AAA(50)-70°C-1hr	16.24
4	BAA(50)-85°C-0.5hr	AAA(50)-70°C-4hr	20.79
5	BAA(50)-85°C-0.5hr	AAA(100)-70°C-1hr	22.13
6	BAA(50)-85°C-0.5hr	AAA(100)-70°C-4hr	23.56

\* BAA: Bacterial alpha-amylase

\* AAA: Acid alpha-amylase

10

## CLAIMS

1. A method of liquefying starch-containing material, wherein the method comprises the steps of  
(a) treating the starch-containing material with a bacterial alpha-amylase at a temperature  
5 around 70-90°C for 15-90 minutes,  
(b) treating the material obtained in step (a) with an alpha-amylase at a temperature between  
60-80°C for 30-90 minutes.
2. The method of claim 1, wherein the starch-containing material is jet-cooking at 90-120°C,  
10 preferably around 105°C, for 1-15 minutes, preferably for 3-10 minute, especially around 5 minutes, before step (a).
3. The method of claims 1 or 2, wherein the starch-containing material is selected from the  
group consisting of: tubers, roots and whole grain; and any combinations of these.  
15
4. The method of any of claims 1-5, wherein the starch-containing material is obtained from cereals.
5. The method of any of the claims 1-5, wherein the starch-containing material is selected from  
20 the group consisting of corn, cob, wheat, barley, rye, milo and potatoes; or any combination of these.
6. The method of claim 5, wherein the starch-containing material is whole grain selected from  
the group consisting of corn, wheat or barley or any combinations of these.  
25
7. The method of any of the claims 1-8, wherein the starch-containing material is whole grain  
and said method comprises a step of milling the whole grain before step (a).
8. The method of any of the claims 1-5, wherein the starch-containing material is obtainable by  
30 a process comprising milling of whole grain.
9. The method of any of the claims 1-8, further comprising prior to step (a) the steps of;  
i) milling of starch-containing material;



ii) forming a slurry comprising the milled material and water.

10. The method of any of the claim 9, wherein the milling step is a dry milling step.

5 11. The method of any of the claims 9-10, wherein the milling step is a wet milling step.

12. The method of any of the claims 1-11, wherein the starch-containing material is a side stream from starch processing.

10 13. The method of claims 1-12, wherein the bacterial alpha-amylase in step (a) is a *Bacillus* alpha-amylase, preferably derived from *Bacillus stearothersophilus* alpha-amylase or a variant with the mutations: I181\*+G182\* especially I181\*+G182\*+N193F.

14. The method of claims 1-13 wherein the alpha-amylase is step (b) is an acid alpha-amylase,  
15 preferably an acid fungal alpha-amylase, preferably derived from *Aspergillus* spp. preferably *Aspergillus niger* or *Aspergillus oryzae*.

15. The method of any of claims 1-14, wherein the acid alpha-amylase is an alpha-amylase having an amino acid sequence which has at least 70% identity to SEQ ID NO:1 preferably at least  
20 75%, 80%, 85% or at least 90%, e.g., at least 95%, 97%, 98%, or at least 99% identity to SEQ ID NO:1.

16. The method of claims 1-15, wherein the acid alpha-amylase is an alpha-amylase having the amino acid sequence set forth in SEQ ID NO:1.

25

17. The method of claims 1-16, wherein the mash obtained after step (b) has a DE value of above 16, preferably above 18, especially above 20, such as a DE value in the range from 16 to 30, preferably in the range from 18 to 25.

30 18. A process of producing ethanol from starch-containing material by fermentation, said process comprises:

- (i) liquefying said starch-containing material as defined in any of claims 1 to 17;
- (ii) saccharifying the liquefied mash obtained;

(iii) fermenting the material obtained in step (ii).

19. The process of any of claim 18, further comprising recovery of the ethanol.

5 20. The process of any of claims 18 or 19, wherein the saccharification and fermentation is carried out as a simultaneous saccharification and fermentation process (SSF process).

21. The process of claim 18, further comprising a step of;

(iv) distillation to obtain the ethanol;

10 wherein the fermentation in step (iii) and the distillation in step (iv) is carried out simultaneously or separately/sequential; optionally followed by one or more process steps for further refinement of the ethanol.

15 22. The process of claim 18, wherein the micro-organism is capable of fermenting sugars to ethanol.

23. The process of claims 18 or 19, wherein the micro-organism is a yeast, such as *Saccharomyces spp.*, preferably *Saccharomyces cerevisiae*.

20 24. The process of any of the claims 18-23, wherein the fermentation is carried out in the presence of a carbohydrate-source generating enzyme..

25 25. The process of claim 24, wherein the carbohydrate-source generating enzyme is a glucoamylase, preferably derived from a strain of *Aspergillus*, preferably *Aspergillus niger* or a strain of *Talaromyces*, especially *Talaromyces emersonii*.

26. The process according to any of claims 18-25, said process comprising the steps of;

1) liquefying starch-containing material in accordance with the liquefaction method of claims 1-17;

30 2) liquefying the material obtained in step 1) in the presence of an alpha-amylase having an amino acid sequence which has at least 70% identity to SEQ ID NO:1; and

3) saccharifying the material obtained in step 2) ; and

4) fermenting to produce ethanol;

wherein the steps 1), 2), 3) and 4) is performed in the order 1), 2), 3), 4) or wherein 4) is performed simultaneously to or following 3).

27. The method of claims 26, wherein the mash obtained after step 2) has a DE value of above  
5 16, preferably above 18, especially above 20, such as a DE value in the range from 16 to 30, preferably in the range from 18 to 25.

## **Abstract**

5 The present invention relates to method of liquefying starch-containing material, wherein the method comprises the steps of (a) treating the starch-containing material with a bacterial alpha-amylase at a temperature around 70-90°C for 15-90 minutes, and (b) treating the material obtained in step (a) with an alpha-amylase at a temperature between 60-80°C for 30-90 minutes. The invention also relates to a process of producing ethanol comprising a liquefaction step carried out according to the liquefaction method of the invention.




Figure 1

## **Application Data Sheet**

### **Application Information**

Application Type::	Provisional
Subject Matter::	Utility
CD-ROM or CD-R::	None
Sequence Submission::	Paper
Computer Readable Form (CRF)::	No
Title::	Liquefaction Process
Attorney Docket Number::	10620.010-US
Request for Early Publication::	No
Request for Non-Publication::	No
Total Drawing Sheets::	0
Small Entity::	No
Petition included::	No
Secrecy Order in Parent Appl.::	No

### **Applicant Information**

Applicant Authority type::	Inventor
Primary Citizenship Country::	India
Status::	Full Capacity
Given Name::	Swapnil
Family Name::	Bhargava
City of Residence::	Raleigh
State or Province of Residence::	North Carolina
Country of Residence::	US

Applicant Authority type::	Inventor
Primary Citizenship Country::	Denmark
Status::	Full Capacity

Given Name:: Henrik  
Family Name:: Bisgard-Frantzen  
City of Residence:: Bagsvaerd  
Country of Residence:: Denmark

Applicant Authority type:: Inventor  
Primary Citizenship Country:: Denmark  
Status:: Full Capacity  
Given Name:: Henrik  
Family Name:: Frisner  
City of Residence:: Vaerlose  
Country of Residence:: Denmark

Applicant Authority type:: Inventor  
Primary Citizenship Country:: Denmark  
Status:: Full Capacity  
Given Name:: Anders  
Family Name:: Vikso-Nielsen  
City of Residence:: Slangerup  
Country of Residence:: Denmark

Applicant Authority type:: Inventor  
Primary Citizenship Country:: US  
Status:: Full Capacity  
Given Name:: Malcolm  
Family Name:: Johal  
City of Residence:: Raleigh  
State or Province of Residence:: North Carolina  
Country of Residence:: US

### **Correspondence Information**

Correspondence Customer Number:: 25908  
Phone Number:: (212) 840-0097  
Fax Number:: (212) 840-0221  
E-Mail address:: Patents-US-NY@novozymes.com

### **Representative Information**

Representative Customer Number:	25908
---------------------------------	-------



# SEQUENCE LISTING

<110> Bhargava, Swapnil  
 Bisgård-Frantzen, Henrik  
 Frisner, Henrik  
 Viksø-Nielsen, Anders  
 Johal, Malcolm

<120> LIQUEFACTION PROCESS

<130> 10620

<160> 1

<170> PatentIn version 3.2

<210> 1

<211> 484

<212> PRT

<213> Aspergillus niger

<220>

<221> misc feature

<223> SEQ ID NO:1

<400> 1

Leu Ser Ala Ala Ser Trp Arg Thr Gln Ser Ile Tyr Phe Leu Leu Thr  
 1 5 10 15

Asp Arg Phe Gly Arg Thr Asp Asn Ser Thr Thr Ala Thr Cys Asn Thr  
 20 25 30

Gly Asn Glu Ile Tyr Cys Gly Gly Ser Trp Gln Gly Ile Ile Asp His  
 35 40 45

Leu Asp Tyr Ile Glu Gly Met Gly Phe Thr Ala Ile Trp Ile Ser Pro  
 50 55 60

Ile Thr Glu Gln Leu Pro Gln Asp Thr Ala Asp Gly Glu Ala Tyr His  
 65 70 75 80

Gly Tyr Trp Gln Gln Lys Ile Tyr Asp Val Asn Ser Asn Phe Gly Thr  
 85 90 95

Ala Asp Asn Leu Lys Ser Leu Ser Asp Ala Leu His Ala Arg Gly Met  
 100 105 110

Tyr Leu Met Val Asp Val Val Pro Asp His Met Gly Tyr Ala Gly Asn

115					120					125					
Gly	Asn	Asp	Val	Asp	Tyr	Ser	Val	Phe	Asp	Pro	Phe	Asp	Ser	Ser	Ser
130						135				140					
Tyr	Phe	His	Pro	Tyr	Cys	Leu	Ile	Thr	Asp	Trp	Asp	Asn	Leu	Thr	Met
145				150						155				160	
Val	Glu	Asp	Cys	Trp	Glu	Gly	Asp	Thr	Ile	Val	Ser	Leu	Pro	Asp	Leu
				165				170						175	
Asp	Thr	Thr	Glu	Thr	Ala	Val	Arg	Thr	Ile	Trp	Tyr	Asp	Trp	Val	Ala
		180						185				190			
Asp	Leu	Val	Ser	Asn	Tyr	Ser	Val	Asp	Gly	Leu	Arg	Ile	Asp	Ser	Val
		195				200						205			
Leu	Glu	Val	Gln	Pro	Asp	Phe	Phe	Pro	Gly	Tyr	Asn	Lys	Ala	Ser	Gly
210						215				220					
Val	Tyr	Cys	Val	Gly	Glu	Ile	Asp	Asn	Gly	Asn	Pro	Ala	Ser	Asp	Cys
225				230						235				240	
Pro	Tyr	Gln	Lys	Val	Leu	Asp	Gly	Val	Leu	Asn	Tyr	Pro	Ile	Tyr	Trp
				245				250						255	
Gln	Leu	Leu	Tyr	Ala	Phe	Glu	Ser	Ser	Ser	Gly	Ser	Ile	Ser	Asn	Leu
		260						265				270			
Tyr	Asn	Met	Ile	Lys	Ser	Val	Ala	Ser	Asp	Cys	Ser	Asp	Pro	Thr	Leu
		275				280						285			
Leu	Gly	Asn	Phe	Ile	Glu	Asn	His	Asp	Asn	Pro	Arg	Phe	Ala	Lys	Tyr
290						295				300					
Thr	Ser	Asp	Tyr	Ser	Gln	Ala	Lys	Asn	Val	Leu	Ser	Tyr	Ile	Phe	Leu
305				310						315				320	
Ser	Asp	Gly	Ile	Pro	Ile	Val	Tyr	Ala	Gly	Glu	Glu	Gln	His	Tyr	Ala
				325				330						335	
Gly	Gly	Lys	Val	Pro	Tyr	Asn	Arg	Glu	Ala	Thr	Trp	Leu	Ser	Gly	Tyr
		340				345						350			

Asp Thr Ser Ala Glu Leu Tyr Thr Trp Ile Ala Thr Thr Asn Ala Ile  
355 360 365

Arg Lys Leu Ala Ile Ala Ala Asp Ser Ala Tyr Ile Thr Tyr Ala Asn  
370 375 380

Asp Ala Phe Tyr Thr Asp Ser Asn Thr Ile Ala Met Ala Lys Gly Thr  
385 390 395 400

Ser Gly Ser Gln Val Ile Thr Val Leu Ser Asn Lys Gly Ser Ser Gly  
405 410 415

Ser Ser Tyr Thr Leu Thr Leu Ser Gly Ser Gly Tyr Thr Ser Gly Thr  
420 425 430

Lys Leu Ile Glu Ala Tyr Thr Cys Thr Ser Val Thr Val Asp Ser Ser  
435 440 445

Gly Asp Ile Pro Val Pro Met Ala Ser Gly Leu Pro Arg Val Leu Leu  
450 455 460

Pro Ala Ser Val Val Asp Ser Ser Ser Leu Cys Gly Gly Ser Gly Arg  
465 470 475 480

Leu Tyr Val Glu